

## DIRECTIONS FOR USE:

**DO NOT** use in greenhouses.

A restricted entry interval (REI) of 1 day is required for workers hand-harvesting and detasseling treated corn.

A restricted entry interval (REI) of 5 days is required for workers hand-harvesting and pruning highbush blueberries.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

**DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

## GENERAL INFORMATION

**DO NOT** graze animals on treated green crops within three days of application of PROPI SUPER 25 ECFungicide.

## PRODUCT INFORMATION

PROPI SUPER 25 EC is a broad-spectrum systemic fungicide for control of a wide range of diseases on certain crops. PROPI SUPER 25 EC fungicide will protect the crop from yield and quality losses due to disease. PROPI SUPER 25 EC may be used in conjunction with higher seeding rates, higher fertilizer inputs, plant growth regulators and other fungicides as required. (See “NOTE” under PRECAUTIONS)

### Diseases Controlled:

CROP	DISEASES CONTROLLED
Barley, Spring	Net Blotch; Spot Blotch; Scald; Powdery Mildew; Septoria Leaf Spot; Leaf Rust; Stem Rust
Canary Seed	Septoria Leaf Mottle
Canola	Blackleg
Corn	Rusts; Northern Corn Leaf Blight; Southern Corn Leaf Blight; Helminthosporium Leaf Spot; Eye Spot; Grey Leaf Spot
Legume Vegetables including Soybeans	Rust ( <i>Phakopsora</i> spp.); Cercospora Leaf Spot ( <i>Cercospora</i> spp.); Powdery mildew ( <i>Microsphaera</i> spp., <i>Erysiphe</i> spp.)
Oats	Septoria Leaf Blotch; Crown Rust
Soybeans grown for seed	Frogeye Leaf Spot ( <i>Cercospora</i> spp.), Aerial Web Blight ( <i>Rhizoctonia solani</i> )
Timothy <sup>1</sup>	Purple eyespot ( <i>Cladosporium phlei</i> )
Wheat, Spring and Winter (including Hard Red, Durum,	Septoria Leaf Spot; Septoria Glume Blotch; Powdery Mildew; Leaf and Stem Rust; Tan Spot; Stripe Rust

Canada Prairie, Soft White)	
Cranberries	Cottonball ( <i>Monilinia oxycocci</i> )
Kentucky Bluegrass Grown for Seed	Powdery Mildew
Lowbush Blueberries	Monilinia Blight (Mummyberry)
Highbush Blueberries	Mummyberry ( <i>Monilinia vaccinii-corymbosi</i> )
Rutabagas	Powdery Mildew
Saskatoon Berry	Entomosporium Leaf and Berry Spot and Saskatoon Juniper Rust
Stone Fruit (Apricots, Nectarines, Peaches, Plums, Sweet and Sour Cherries)	Brown Rot, Blossom Blight and Fruit Brown Rot, Cherry Leaf Spot ( <i>Blumeriella jaapii</i> – sweet and sour cherries)
Plums and Sour Cherries	Black Knot ( <i>Apiosporina morbosa</i> ) – Suppression ONLY
Western Red Cedar	Keithia Foliar Blight
Asparagus	Rusts ( <i>Puccinia asparagi</i> )
Caneberries	Yellow Rust
Strawberries	Common leaf spot ( <i>Mycosphaerella fragariae</i> )

<sup>1</sup> FOR SALE FOR USE ON TIMOTHY HAY IN THE PRAIRIE PROVINCES ONLY.

**Factors Affecting PROPI SUPER 25 EC Performance:** PROPI SUPER 25 EC should be applied as a preventative disease control measure. Established diseases are more difficult to control and may have already reduced crop vigour.

If rainfall occurs within one hour of application, reapplication is necessary.

#### **Field Sprayer Application:**

**DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

#### **Airblast application:**

**DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

#### **AERIAL APPLICATION:**

**DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotor span. High humidity and low temperatures (10 - 20°C) allow for a better deposition of spray droplets.

## SPRAYER AND APPLICATION INFORMATION:

The performance of this product depends on correct application. Follow the guidelines given below for optimal application of PROPI SUPER 25 EC.

### Sprayer Information:

	GROUND APPLICATION	AERIAL APPLICATION
Spray Volume	Minimum 200 L of water per hectare	40 - 50 L of water per hectare
Spray Pressure	200 - 300 kPa	100 - 200 kPa
Nozzle Type	110° Flat Fan (XR11004, 4110-20)	Flat Fan 6510-6515 or Hollow Cone (D8-45)
Droplet Size	Medium Spray (300 - 400 microns VMD)	Medium Spray (350 - 400 microns VMD)
Ground Speed	10 km/h	
Nozzle Angle	90° (straight down)	90° (straight down)
Boom Height	40-50 cm above the crop canopy	2-3 m above the crop canopy

### Ground Application - Mixing and Spraying Instructions:

1. Spray equipment should be thoroughly flushed with clean water before mixing PROPI SUPER 25 EC.
2. Fill spray tank 1/2 full with clean water. Engage gentle agitation.
3. Add the required amount of PROPI SUPER 25 EC and agitate thoroughly.
4. Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tank mix partner.
5. Complete filling the spray tank with water, maintaining agitation during mixing and spraying operations.
6. Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

### Aerial Application - Mixing and Spraying Instructions:

1. Spray equipment should be thoroughly flushed with clean water before mixing PROPI SUPER 25 EC.
2. Fill premix tank 1/2 full with clean water. Engage gentle agitation.
3. Add the required amount of PROPI SUPER 25 EC and agitate thoroughly.
4. Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tank mix partner.
5. Complete filling the premix tank with water.
6. Maintain gentle agitation during mixing.
7. Transfer the premix contents into the aircraft spray tank.
8. Maintain sufficient agitation during the mixing and spraying operation to ensure PROPI SUPER 25 EC remains in suspension.
9. Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for a specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

**AERIAL APPLICATION USE PRECAUTIONS:**

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift; therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as a shelter-belt) or aquatic habitat.

**AERIAL APPLICATION OPERATOR PRECAUTIONS:**

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the **ground application section**.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

**Buffer zones:**

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas,

shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method Of Application	Crop		Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitats of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer*	Beans, peas, soybeans, chickpeas, corn, wheat, oats, rye, triticale canary seed, canola, barley, rutabagas, cranberries, strawberries, asparagus, Kentucky bluegrass, Western cedar		1	0	1	1	1
Aerial	beans, corn, oats, wheat, barley, triticale (seed prod.), blueberries, Kentucky bluegrass (seed prod.)	Fixed Wing	1	0	3	1	20
		Rotary Wing	1	0	1	1	20
Airblast	Cherries	Early Growth Stage	5	0	10	3	10
		Late Growth Stage	2	0	4	2	4
	Blueberries, apricots, nectarines, peaches, plums, Saskatoon berries, blackberries, loganberries, raspberries, other berries	Early Growth Stage	4	0	5	2	5
		Late Growth Stage	2	0	3	1	3

\*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled **bufferzone** can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

**PRODUCT SPECIFIC PRECAUTIONS:**

Read and understand the entire label before opening this product. If you have questions call the manufacturer at 1-888-760-0187 or obtain technical advice from the distributor or your provincial agricultural representative.

**Fertilizer - Mixing and Spraying Instructions:**

If desired, small amounts of nitrogen may be applied with PROPI SUPER 25 EC. The appropriate amount of urea can be dissolved in water and added to the spray tank before adding PROPI SUPER 25 EC. The rate of actual nitrogen must not exceed 10 kg/ha.

CAUTION: Excessive nitrogen concentrations may injure the crop.

NOTE: DO NOT add nitrogen when tank-mixing PROPI SUPER 25 EC with an herbicide.

**WHEAT, BARLEY AND OATS INSTRUCTIONS FOR USE:**

Apply PROPI SUPER 25 EC at the very early stages of disease. This could occur anytime during tillering or stem elongation. Typically, an application from the beginning of stem elongation up to flag leaf emergence is required.

PROPI SUPER 25 EC lasts about three weeks in the plant. If conditions favourable to disease continue after this length of time, another application will be necessary to maintain control. The second spray is usually applied at the time of head emergence. In most cases, this second application is essential to maintain control of the Septoria disease complex.

LAST APPLICATION MUST BE MADE AT LEAST 45 DAYS BEFORE HARVEST (45 day PHI).

<b>CROP/DISEASES</b>	<b>WHEAT: Septoria Leaf Spot, Tan Spot SPRING BARLEY: Net Blotch</b>
<b>RATE/HA</b>	250 - 500 mL
<b>EARLY Application</b>	At G.S. 12-23 (as early as the two leaf stage). For early season disease suppression, use the lower rate for suppression under normal field conditions. Use the higher rate for control if there is a history of high disease pressures in the field and/or field conditions favour disease development.
<b>LATER Application</b>	At the first sign of disease (G.S. 29-37) or before head is half emerged (G.S. 49-55). Apply only the high rate on any application from G.S. 29-55.
<b>MAXIMUM NUMBER OF</b>	2

<b>APPLICATIONS PER SEASON</b>	
<b>CROP/DISEASES</b>	WHEAT: <b>Septoria Leaf Spot and Glume Blotch, Powdery Mildew, Leaf and Stem Rust, Tan Spot, Stripe Rust</b> SPRING BARLEY: <b>Net Blotch, Spot Blotch, Scald, Powdery Mildew,</b> <b>Septoria Leaf Spot, Leaf and Stem Rust</b> OATS: <b>Septoria Leaf Blotch, Crown Rust</b>
<b>RATE/HA</b>	500 mL
<b>EARLY Application</b>	At the first sign of disease, usually at the beginning of stem elongation (G.S. 29-37).
<b>LATER Application</b>	Before head is half emerged (G.S. 49-55).
<b>MAXIMUM NUMBER OF APPLICATIONS PER SEASON</b>	2

#### **HERBICIDE TANK-MIXING - WHEAT & BARLEY:**

PROPI SUPER 25 EC can be tank-mixed with ONLY ONE of these herbicides:

2,4-D Amine

MCPA Amine

Buctril® M

Pardner®

Mextrol® 450

Approve®

HORIZON® 240EC Herbicide (Wheat Only)

AXIAL® 100EC Herbicide (Spring Wheat and Spring Barley Only) (FOR USE ONLY IN THE PRAIRIE AND PEACE RIVER, OKANAGAN AND CRESTON FLATS REGIONS OF BRITISH COLUMBIA)

BROADBAND® Herbicide (Spring Wheat (excluding durum) and Barley) (FOR USE IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA ONLY)

#### **Tank-mixing Precautions:**

1. DO NOT tank-mix PROPI SUPER 25 EC Fungicide with herbicides for application onto Oats.
2. Weeds and crops must be at the correct stage of growth as specified in both the PROPI SUPER 25 EC Fungicide label and the tank mix partner label.
3. 2,4-D and MCPA formulations may be applied either by ground application or aerial application; tank-mixtures of PROPI SUPER 25 EC Fungicide and Buctril M, Mextrol 450, Approve, Pardner, or AXIAL 100EC Herbicide can only be applied by ground application.
4. For fields treated with AXIAL 100EC Herbicide, no crop may be seeded until the following year. There are no crop rotation limitations the year following application of AXIAL 100EC Herbicide.

5. Tank mixing with the herbicide AXIAL 100EC will slightly decrease PROPI SUPER 25 EC efficacy.
6. Consult the label of the herbicide partner for a list of weeds controlled, directions for use and precautions.
7. When tank mixing, adhere to the most restrictive label limitations and precautions.
8. Compatibility should always be confirmed by premixing small proportional quantities of water, PROPI SUPER 25 EC Fungicide, and the tank-mix partner in advance.

**NOTE:**

- DO NOT graze animals on treated green crops within three days of application of PROPI SUPER 25 EC Fungicide.
- DO NOT feed straw from crops treated with herbicide tank mixes to livestock.

**HERBICIDE TANK-MIXING – WHEAT ONLY:**

FOR USE ONLY IN THE PRAIRIE AND PEACE RIVER, OKANAGAN AND CRESTON FLATS REGIONS OF BRITISH COLUMBIA.

PROPI SUPER 25 EC can be tank-mixed with HORIZON 240 EC Herbicide for disease and grassy weed control.

**Tank-mixing Precautions:**

1. Do not apply by air.
2. Consult the label of HORIZON 240 EC Herbicide for a list of weeds controlled, directions for use and precautions.
3. Apply prior to emergence of the 4th tiller (herbicide timing).

**TIMOTHY - FOR SALE FOR USE IN THE PRAIRIE PROVINCES ONLY**

**INSTRUCTIONS FOR USE:**

Apply PROPI SUPER 25 EC at the very early stages of disease. If conditions favourable to disease continue after this length of time, another application will be necessary to maintain control. DO NOT APPLY BY AIR.

LAST APPLICATION MUST BE MADE PRIOR TO 14 DAYS BEFORE HARVEST (14 day PHI).

<b>DISEASE</b>	Purple eyespot
<b>RATE/HA</b>	500 ml
<b>EARLY Application</b>	At the first sign of disease, usually at the beginning of flowering (G.S. 59-61).
<b>LATER Application</b>	Full flowering (G.S. 65-73).
<b>Minimum Interval between Applications</b>	14 days
<b>Maximal Seasonal Application Rate</b>	2 applications at 500 mL/ha (1.0 L/ha in a season).

**CANOLA**

**INSTRUCTIONS FOR USE:**



PROPI SUPER 25 EC will control blackleg and enhance yield potential during the early stages of canola growth. The disease may reappear later in the season, but with minimal effect on yield. LAST APPLICATION MUST BE MADE PRIOR TO 60 DAYS BEFORE HARVEST (60 day PHI).

DISEASE	RATE/HA	REMARKS/TIMING
Blackleg	500 mL	<b>Apply during the rosette stage;</b> between 2 <sup>nd</sup> true leaf and bolting.
		Seedling <b>Rosette</b> Bud (Bolted) Stage 1      Stage 2      Stage 3

### TANK-MIXING – SEED CORN, FIELD CORN and SWEET CORN

PROPI SUPER 25 EC can be tank-mixed with the following partner:

MATADOR<sup>®</sup> 120EC Insecticide

WARRIOR<sup>®</sup> Insecticide

Ripcord<sup>®</sup>

#### Tank-mixing Precautions:

- DO NOT APPLY THE TANK MIX WITH WARRIOR INSECTICIDE BY AIR.
- The tank mix of PROPI SUPER 25 EC + MATADOR 120EC Insecticide or PROPI SUPER 25 EC + Ripcord can be applied by air and ground. Use 40 L of water per hectare when applying by air.
- Insects and crops must be at the correct stage as specified on the PROPI SUPER 25 EC as well as MATADOR 120EC, WARRIOR Insecticide and Ripcord (PCP 15738) labels. Follow the directions for use and precautions on all labels.
- DO NOT apply within 14 days of harvest (14 day PHI) when using these tank-mixes on field and sweet corn.
- Compatibility should always be confirmed by premixing small proportional quantities of water, PROPI SUPER 25 EC, and the tank-mix partner in advance.

### TANKMIXES SEED CORN

DISEASES	Rusts
RATE/HA PROPI SUPER 25 ECalone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of PROPI SUPER 25 EC Fungicide when rust pustules first appear. Under severe disease pressure, make a second application 14 days after. Under severe disease pressure, make a third application 14 days after. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Northern Corn Leaf Blight Southern Corn Leaf Blight

	Helminthosporium Leaf Spot
RATE/HA PROPI SUPER 25 ECalone	250 - 500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 250 - 500 mL/ha of PROPI SUPER 25 EC Fungicide when disease first appears. Use the 250 mL rate if disease pressure is low. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Eye Spot Grey Leaf Spot
RATE/HA PROPI SUPER 25 EC alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL of PROPI SUPER 25 EC Fungicide per hectare when disease first appears. DO NOT apply the tank mix with WARRIOR Insecticide by air.

#### FIELD CORN AND SWEET CORN

DISEASES	Rusts
RATE/HA PROPI SUPER 25 EC alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of PROPI SUPER 25 EC Fungicide when rust pustules first appear. Under severe disease pressure, make a second application 14 days after. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Northern Corn Leaf Blight Southern Corn Leaf Blight Helminthosporium Leaf Spot
RATE/HA PROPI SUPER 25 EC alone	250 - 500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 250 - 500 mL/ha of PROPI SUPER 25 EC Fungicide when disease first appears. Use the 250 mL rate if disease pressure is low.

	DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Eye Spot Grey Leaf Spot
RATE/HA PROPI SUPER 25 EC alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL of PROPI SUPER 25 EC Fungicide per hectare when disease first appears. DO NOT apply the tank mix with WARRIOR Insecticide by air.

### LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS

#### Soybeans

**Edible-podded legume vegetables** - Any succulent cultivar of edible podded bean (*Phaseolus* spp.) and any succulent cultivar of edible-podded pea (*Pisum* spp.): Bean (*Phaseolus* spp.) (includes runner bean, snap bean, wax bean); bean (*Vigna* spp.) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean); jackbean; pea (*Pisum* spp.) (includes dwarf pea, edible-podded pea, snow pea, sugar snap pea); pigeon pea; soybean (immature seed); and sword bean.

**Succulent shelled pea and bean** - Any succulent shelled cultivar of bean (*Phaseolus* spp.) and garden pea (*Pisum* spp.): Bean (*Phaseolus* spp.) (includes lima bean, green bean); broad bean (succulent); bean (*Vigna* spp.) (includes black-eyed pea, cowpea, southern pea); pea (*Pisum* spp.) (includes English pea, garden pea, green pea); and pigeon pea.

**Dried shelled pea and bean** - Any dried cultivar of bean (*Phaseolus* spp.) and dried cultivar of pea (*Pisum* spp.): Dried cultivars of bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, and sweet white lupin); (*Vigna* spp.) (includes adzuki beans); (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean, mung bean, rice bean, southern bean, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil; pea (*Pisum* spp.) (includes field pea); and pigeon pea.

### LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS

DISEASE	Asian (Soybean) Rust ( <i>Phakopsora pachyrhizi</i> )
CROPS	All Crop Group 6 legume vegetables listed above, and soybean
RATE/HA	500 – 756 mL
REMARKS	For control of Asian Rust, apply the lower rate under conditions of low disease pressures. Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development. APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF PROPI SUPER 25 EC. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry

	legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).
DISEASE	Powdery Mildew ( <i>Microsphaera diffusa</i> , <i>Erysiphe pisi</i> , <i>E. polygoni</i> )
CROPS	All Crop Group 6 legume vegetables listed above, and soybean
RATE/HA	500 mL
REMARKS	Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development. APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF PROPI SUPER 25 EC. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).
DISEASE	Cercospora Leaf Spot ( <i>Cercospora kikuchi</i> )
CROPS	Soybean only
RATE/HA	500 mL
REMARKS	Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development. APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF PROPI SUPER 25 EC. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).

**TANK MIX - PROPI SUPER 25 EC FUNGICIDE with QUADRIS® Flowable Fungicide  
LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS**

DISEASE	Asian (Soybean) Rust ( <i>Phakopsora pachyrhizi</i> )
CROPS	All Crop Group 6A, B, C legume vegetables listed above, and soybeans
RATE/HA	0.5 - 0.75 L/ha of PROPI SUPER 25 EC Fungicide + 0.3 - 0.45 L/ha of QUADRIS Flowable Fungicide
REMARKS	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressures. A second application at 14 days interval may be needed if conditions persist. It is important to protect the developing pod of soybean and podded legume vegetables. Good spray coverage and canopy penetration are important for best results. Apply in a minimum of 45 L of water per hectare.
Restrictions: DO NOT APPLY MORE THAN 2 APPLICATIONS OF PROPI SUPER 25 EC FUNGICIDE PER SEASON. DO NOT APPLY MORE THAN 2 APPLICATIONS OF QUADRIS FLOWABLE FUNGICIDE PER SEASON. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT	

apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables). DO NOT make more than one application to soybean hay and dry pea hay. DO NOT apply within 14 days of harvest (14 day PHI) of soybean hay and dry pea hay. DO NOT feed dried pea vines to livestock. Not all members of the legume vegetable group have been tested for efficacy and phytotoxicity at the recommended label rates and this tank mix should be used at the discretion of the user. The tank mix of PROPI SUPER 25 EC Fungicide plus QUADRIS Flowable Fungicide may be applied by air or ground application equipment.

**TANK MIX - PROPI SUPER 25 EC FUNGICIDE with MATADOR 120EC Insecticide or WARRIOR Insecticide  
LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS**

PROPI SUPER 25 EC Fungicide can be tank mixed with MATADOR 120EC Insecticide or WARRIOR Insecticide for foliar disease and insect control. Apply PROPI SUPER 25 EC Fungicide at a rate of 250 - 756 mL/ha in a tank mix with MATADOR 120EC Insecticide or WARRIOR Insecticide at a rate of 83 - 233 mL/ha for control of soybean aphid on soybean and at a rate of 83 mL/ha for the rest of the crops in Crop Group 6. Refer to both the PROPI SUPER 25 EC Fungicide, MATADOR 120EC Insecticide, or WARRIOR Insecticide labels for diseases and insects controlled, specific application instructions, and precautions. Pests and crops must be at the correct stage as specified on the PROPI SUPER 25 EC Fungicide, MATADOR 120EC Insecticide, or WARRIOR Insecticide labels. DO NOT apply more than 2 applications per season of this tank mix. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables). DO NOT graze or harvest treated forage, straw or hay for livestock feed. Not all members of the legume vegetable group have been tested for efficacy and phytotoxicity at the recommended label rates, and should be used at the discretion of the user. This tank mix can be applied by ground application equipment only. DO NOT APPLY THE TANK MIX WITH MATADOR 120EC Insecticide or WARRIOR Insecticide BY AIR.

**TANK MIXES - PROPI SUPER 25 EC FUNGICIDE with QUADRIS Flowable Fungicide  
CEREALS – CROP GROUP 15**

CROPS	Barley, Oats
DISEASE	Barley net blotch ( <i>Pyrenophora teres</i> )
PRODUCT RATE (L/ha)	0.5 L/ha of PROPI SUPER 25 EC Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Barley, Rye
DISEASE	Barley scald ( <i>Rhynchosporium secalis</i> )
PRODUCT RATE (L/ha)	0.5 L/ha of PROPI SUPER 25 EC Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide

APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Wheat, Barley, Rye, Oats, Triticale
DISEASE	Septoria leaf spot ( <i>Septoria</i> sp.)
PRODUCT RATE (L/ha)	0.5 L/ha of PROPI SUPER 25 EC Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Wheat, Barley, Rye, Triticale
DISEASE	Tan spot ( <i>Pyrenophora tritici-repentis</i> )
PRODUCT RATE (L/ha)	0.5 L/ha of PROPI SUPER 25 EC Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Barley
DISEASE	Barley leaf rust ( <i>Puccinia hordei</i> )
PRODUCT RATE (L/ha)	0.5 L/ha of PROPI SUPER 25 EC Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Winter Wheat, Spring Wheat, Barley
DISEASE	Stripe rust ( <i>Puccinia striiformis</i> )
PRODUCT RATE (L/ha)	0.4 - 0.5 L/ha of PROPI SUPER 25 EC Fungicide + 0.2 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Winter Wheat, Spring Wheat
DISEASE	Wheat leaf rust ( <i>Puccinia triticina</i> )
PRODUCT RATE (L/ha)	0.4 - 0.5 L/ha of PROPI SUPER 25 EC Fungicide + 0.2 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.

**Specific Use Restrictions:** DO NOT make more than one application per season of this tank mixture. An additional application of PROPI SUPER 25 EC Fungicide can be made, if required. Refer to the PROPI SUPER 25 EC Fungicide label for details of the rate and timing. A total of two applications of PROPI SUPER 25 EC Fungicide should be applied per season either in a tank mix with QUADRIS Flowable Fungicide or alone. DO NOT apply within 45 days of harvest (45 day PHI) for grain and straw. DO NOT harvest wheat for forage. DO NOT graze or feed livestock treated forage or cut green crop for hay or silage. The tank mix of QUADRIS Flowable Fungicide and PROPI SUPER 25 EC Fungicide may be applied with ground or air equipment.

**GROUND APPLICATION:** Apply specified rates in a minimum of 100 L of water per hectare.

**AERIAL APPLICATION:** Apply specified rates in a minimum of 45 L of water per hectare.

**TANK MIX - PROPI SUPER 25 EC FUNGICIDE with QUADRIS Flowable Fungicide  
CORN**

CROPS	Field Corn, Sweet Corn (including Seed Production), Popcorn (including Seed Production)
DISEASE	Rust ( <i>Puccinia sorghi</i> ) Northern Corn Leaf Blight ( <i>Setosphaeria turcicum</i> ) Southern Corn Leaf Blight ( <i>Cochliobolus heterostrophus</i> ) Eye Spot ( <i>Aureobasidium zeae</i> ) Grey Leaf spot ( <i>Cercospora zeae-maydis</i> )
PRODUCT RATE (L/ha)	0.5 L/ha of PROPI SUPER 25 EC Fungicide + 0.225 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Make first application at the first sign of disease, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development. Good spray coverage and canopy penetration are important for best results. Apply in a minimum of 45 L of water per hectare. Use the low rate of QUADRIS under low to moderate disease pressure. Use the high rate of QUADRIS only under conditions of severe disease pressure.
<p><b>Specific Use Restrictions:</b> A maximum of two applications of QUADRIS Flowable Fungicide and a maximum of two applications of PROPI SUPER 25 EC Fungicide should be applied per season either as a tank mix or as products applied alone. DO NOT apply to field corn and field corn grown for seed after brown silk.</p> <p>DO NOT apply within 30 days (30 day PHI) of harvest for forage. DO NOT apply to sweet corn within 14 days of harvest (14 day PHI).</p> <p>Not all of these cereal crops have been tested for efficacy and phytotoxicity at the recommended label rates and the tank mix of QUADRIS Flowable Fungicide and PROPI SUPER 25 EC Fungicide should be used at the discretion of the user. The tank mix of QUADRIS Flowable Fungicide and PROPI SUPER 25 EC Fungicide may be applied with ground or air equipment.</p> <p><b>GROUND APPLICATION:</b> Apply specified rates in a minimum of 100 L of water per hectare.</p> <p><b>AERIAL APPLICATION:</b> Apply specified rates in a minimum of 45 L of water per hectare.</p>	

RUTABAGAS		
DISEASE	RATE/HA	REMARKS

PowderyMilde w	400 mL	Make two applications per season with the first application at 50 days after planting and the 2nd application 20 days later. Apply to vegetative foliage. Use 200 L of water per hectare. LAST APPLICATION MUST BE MADE AT LEAST 21 DAYS BEFORE HARVEST. Do not graze livestock on treated crop.
-------------------	--------	--

WESTERN RED CEDAR		
DISEASE	RATE/HA	REMARKS
Keithia Foliar Blight	500 mL	Apply using ground application equipment every four weeks. Make a maximum of 6 applications per year. Apply in a volume of 1000 L of water per hectare. Do not graze livestock on treated crop.

HIGH BUSH BLUEBERRY		
DISEASE	RATE/HA	REMARKS
Mummyberry ( <i>Monilinia vaccinii-corymbosi</i> )	500 mL	Apply first application at or near flower bud swelling; make a second application at leaf bud swelling. Apply by ground only, making no more than two applications per year. In BC only, apply by ground, a third application at pink bloom and a fourth application 7 to 10 days later at early bloom, making no more than 4 applications per year. Use a minimum of 200 L of water per hectare. LAST APPLICATION MUST BE MADE PRIOR TO 60 DAYS BEFORE HARVEST (60 day PHI). Do not graze livestock on treated crop.

LOW BUSH BLUEBERRY		
DISEASE	RATE/HA	REMARKS
Monilinia Blight	500 mL	Apply first application when flower bud scales first appear and make a second application 10 days later. Use ground application or aerial application equipment, making no more than two applications per year. Use a minimum of 200 L of water per hectare if applying by ground equipment; use 40-50 L of water per hectare if applying by air. LAST APPLICATION MUST BE MADE PRIOR TO 60 DAYS BEFORE HARVEST (60 day PHI). Do not graze livestock on treated crop.

CRANBERRY		
DISEASE	RATE/HA	REMARKS



Cottonball( <i>Monilinia</i> <i>xyccoci</i> )	500 mL	Apply the first application at leaf bud break. Make a second application 10 -14 days later, a third application at early bloom, and a fourth application 10 - 14 days after the third application. Make no more than four applications per year. LAST APPLICATION MUST BE MADE PRIOR TO 45 DAYS BEFORE HARVEST(45 day PHI). Do not graze livestock on treated crop.
---	--------	--

<b>SASKATOONBERRY</b>		
<b>DISEASE</b>	<b>RATE/HA</b>	<b>REMARKS</b>
Entomosporium Leafand BerrySpot, Saskatoon JuniperRust	500 mL	As a foliar spray, apply up to three applications per season. The first application to occur at “white tip”, the second application at “petal fall”, and the third application at “green fruit”. Use a minimum of 200 L of water per hectare by ground, applying to runoff. LAST APPLICATION MUST BE MADE PRIOR TO 38 DAYS BEFORE HARVEST (38 day PHI). Do not graze livestock on treated crop.

#### **PEACHES, NECTARINES, PLUMS, SWEET & SOUR CHERRIES, APRICOTS**

DISEASE	Brown Rot, Blossom Blight
RATE/HA	500 mL
REMARKS	Apply 500 mL of PROPI SUPER 25 EC in 500 L of water per hectare. Make 1st application at early bloom with a 2nd application at 50 - 75% bloom. If disease conditions persist, make a 3rd application at petal fall. Do not graze livestock on treated crop.
DISEASE	Fruit Brown Rot
RATE/HA	500 mL
REMARKS	Apply no more than 2 applications in the 3 weeks prior to harvest. Apply 500 mL of product in 500 L of water per hectare. Do not graze livestock on treated crop.
DISEASE	Suppression of Black Knot (Plums and Sour Cherries ONLY)
RATE/HA	500 mL
REMARKS	Apply 500 mL of PROPI SUPER 25 EC in a minimum of 500 L of water per hectare by ground. Make 1st application at early bloom with 2nd application at 50 - 75% bloom. If disease conditions persist, make 3rd application at petal fall. Do not graze livestock on treated crop.

#### **PEACHES, NECTARINES, PLUMS, SWEET & SOUR CHERRIES, APRICOTS**

DO NOT APPLY WITHIN 3 DAYS OF HARVEST (3 day PHI). DO NOT REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION (3 day REI). IF REQUIRED, INDIVIDUALS MAY REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION FOR SHORT TERM TASKS WHICH DO NOT REQUIRE HAND LABOUR IF AT LEAST 4 HOURS HAS PASSED SINCE APPLICATION AND PROVIDED LONG PANTS, LONG SLEEVED SHIRTS, AND CHEMICAL RESISTANT GLOVES ARE WORN.

FOR SOUR CHERRIES:

a) It is recommended that no more than 2 consecutive applications of PROPI SUPER 25 EC be made before switching to another fungicide with a different mode of action according to disease management practices.

b) Apply a MAXIMUM of 5 APPLICATIONS PER SEASON of PROPI SUPER 25 EC.

#### KENTUCKY BLUEGRASS FOR SEED PRODUCTION

DISEASE	RATE/HA	REMARKS
Powdery Mildew	500 mL	Apply as a foliar spray. Make no more than 2 applications per crop year with the 1st application at pre-heading and the 2nd at 50%-100% heading. Apply in 200-300L/ha of water by ground or 40-50L/ha of water by air. <b>Do not graze livestock on treated crop.</b>

#### ASPARAGUS

DISEASE	RATE/HA	REMARKS
Rust ( <i>Puccinia asparagi</i> )	250mL	Apply PROPI SUPER 25 EC to asparagus ferns in Ontario and Quebec only. Once harvest is complete, make the first application of PROPI SUPER 25 EC as soon as fern growth begins, followed by applications at 14 - 21 day intervals. For new, non-harvested plantings, apply PROPI SUPER 25 EC when first sign of rust is visible, followed by applications at 14 - 21 day intervals. Apply by ground only, making no more than three applications per year. Use a minimum of 370 L of water per hectare. <b>LAST APPLICATION MUST BE MADE PRIOR TO 8 MONTHS BEFORE HARVEST (8 month PHI). Do not graze livestock on treated crop.</b>

#### NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Sharda Cropchem Limited and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Sharda Cropchem Limited itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance

arising, and agree to hold Sharda Cropchem Limited harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.	
<b>SOYBEANS GROWN FOR SEED (Minor Use)</b>	
DISEASES	Frogeye Leaf Spot ( <i>Cercospora</i> spp.) Aerial Web Blight ( <i>Rhizoctonia solani</i> )
RATE/HA	500 - 760 mL
REMARKS	Apply using ground application equipment only, when disease first appears. Under severe disease pressure, make a second application 14 days after the first application. DO NOT harvest soybeans within 30 days of the last application (30 day PHI). For use only on soybeans grown for seed. Harvested soybean seed should not be used for human food or animal feed.
<b>CANARY SEED (Minor Use)</b>	
DISEASE	Septoria Leaf Mottle
RATE/HA	500 mL
REMARKS	For the suppression of Septoria leaf mottle. Make one application at emergence of flag leaf. Ground application only. Apply in 200 L water/ha.
<b>DRY EDIBLE BEANS (Minor Use)</b>	
DISEASE	Rust
RATE/HA	500 mL
REMARKS	Apply 500 mL of PROPI SUPER 25 EC in minimum of 200 L water/ha by ground application or in 40 - 50 L water/ha by aerial application at the first detection of disease in the field and a second application 14 to 21 days later. APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF PROPI SUPER 25 EC TO DRYEDIBLE BEANS. DO NOT APPLY WITHIN 28 DAYS OF HARVEST (28 day PHI).
<b>SWEET AND SOUR CHERRIES (Minor Use)</b>	
DISEASE	Cherry Leaf Spot ( <i>Blumeriella jaapii</i> )
RATE/HA	500 mL
REMARKS	Maximum of three applications per season. Ground application by airblast sprayer. Retreatment interval 7 - 10 days. DO NOT apply within 3 days of harvest (3 day PHI). It is recommended that no more than 2 consecutive applications of PROPI SUPER 25 EC be made before switching to another fungicide with a different mode of action according to disease management practices. Do not graze livestock on treated crop.
<b>CANEBERRIES (Crop Group 13A; cultivars and/or varieties of red and black raspberry, loganberry, and blackberry) (Minor Use)</b>	
DISEASE	Yellow Rust
RATE/HA	500 mL
REMARKS	Apply 500 mL product per hectare in 500 L water by ground application at first detection of disease in the field and then 14 days later. Apply a maximum of two applications per season. DO NOT apply within 30 days of harvest (30 day PHI). Do not graze livestock on treated crop.
<b>STRAWBERRIES (Minor Use)</b>	

DISEASE	Common Leaf Spot ( <i>Mycosphaerella fragariae</i> )
RATE/HA	500 mL
REMARKS	Apply 500 mL of PROPI SUPER 25 EC per hectare by ground in enough water to ensure thorough coverage. Make first application when disease levels are no more than 5%. Apply at 10day intervals for control of leaf spot. It is recommended that no more than 2 consecutive applications of PROPI SUPER 25 EC be made before switching to another fungicide with a different mode of action, according to disease management practices. APPLY A MAXIMUM OF 4 APPLICATIONS PER SEASON OF PROPI SUPER 25 EC TO STRAWBERRIES. DO NOT APPLY WITHIN 1 DAY OF HARVEST (1 day PHI). Do not graze livestock on treated crop.

### RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, PROPI SUPER 25 EC contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to PROPI SUPER 25 EC and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

#### To delay fungicide resistance:

Where possible, rotate the use of PROPI SUPER 25 EC or other Group 3 fungicides with different groups that control the same pathogens.

Use tank mixtures with fungicide from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for resistance development. Notify Sharda CropChem Limited if reduced sensitivity of the pathogen to PROPI SUPER 25 EC is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact company representatives at 1-888-760-0187.